



Informed Consent

Rhinoplasty Surgery



INSTRUCTIONS

This is an informed consent document that has been prepared to help inform you about rhinoplasty surgery, its risks, as well as alternative treatment(s).

It is important that you read this information carefully and completely. Please initial each page, indicating that you have read the page and sign the consent for surgery as proposed by your plastic surgeon and agreed upon by you.

GENERAL INFORMATION

Surgery of the nose (rhinoplasty) is an operation frequently performed by plastic surgeons. This surgical procedure can produce changes in the appearance, structure, and function of the nose. Rhinoplasty can reduce or increase the size of the nose, change the shape of the tip, narrow the width of the nostrils, or change the angle between the nose and the upper lip. This operation can help correct birth defects, nasal injuries, and help relieve some breathing problems.

There is not a universal type of rhinoplasty surgery that will meet the needs of every patient. Rhinoplasty surgery is customized for each patient depending on his or her needs. Incisions may be made within the nose or concealed in inconspicuous locations on the nose. In some situations, cartilage grafts taken from within the nose or from other areas of the body may be recommended to help reshape the structure of the nose. Internal nasal surgery to improve nasal breathing can be performed at the time of the rhinoplasty.

The best candidates for this type of surgery are individuals who are looking for improvement, not perfection, in the appearance of their nose. In addition to realistic expectations, good health and psychological stability are important qualities for patients considering rhinoplasty surgery. Rhinoplasty can be performed in conjunction with other surgeries.

ALTERNATIVE TREATMENTS

Alternative forms of management include not undergoing the rhinoplasty surgery. Certain internal nasal airway disorders may not require surgery on the exterior of the nose. Some nasal contour problems may be treated with temporary injectable fillers. Risks and potential complications are associated with alternative surgical forms of treatment.

INHERENT RISKS OF RHINOPLASTY SURGERY

Every surgical procedure involves a certain amount of risk and it is important that you understand these risks and the possible complications associated with them. In addition, every procedure has limitations. An individual's choice to undergo a surgical procedure is based on the comparison of the risk to the potential benefit. Although the majority of patients do not experience these complications, you should discuss each of them with your plastic surgeon to make sure you understand all the possible consequences of rhinoplasty.

PROCEED TO NEXT PAGE

Patient Name:

Patient Initials: _____

SPECIFIC RISKS OF RHINOPLASTY SURGERY

Damage to Donor Sites:

Occasionally, the procedure will involve using cartilage or bone grafts taken from another site on the body. This adds risks that your surgeon will discuss with you.

Implants:

An implant may be used in some rhinoplasty operations. Implants may become infected or exposed, requiring removal. This may occur at any time in the future.

Delayed Healing:

Fracture disruption or delayed wound healing is possible. Some areas of the nose may not heal normally and may take a long time to heal. Areas of skin may die. This may require frequent dressing changes or further surgery to remove the non-healed tissue.

Nasal Septal Perforation:

Infrequently, a hole in the nasal septum will develop. The occurrence of this is rare. Additional surgical treatment may be necessary to repair the nasal septum. In some cases, it may be impossible to correct this complication.

Nasal Airway Alterations:

Changes may occur after a rhinoplasty or septoplasty operation that may interfere with the normal passage of air through the nose.

Substance Abuse Disorders:

Individuals with substance abuse problems that involve the inhalation of vasoconstrictive drugs such as cocaine are at risk for major complications including poor healing and nasal septal perforation.

PROCEED TO NEXT PAGE

Patient Name:

Patient Initials: _____

GENERAL RISKS OF SURGERY

Healing Issues:

Certain medical conditions, dietary supplements, and medication may delay and interfere with healing. Patients with massive weight loss may have a healing delay that could result in the incisions coming apart, infections, and tissue changes that would require additional medical care, surgeries, and prolonged hospitalizations. Patients with diabetes, or those taking medications (e.g., steroids) on an extended basis may have prolonged healing issues. Smoking will cause a delay in the healing process, often resulting in the need for additional surgeries. There are general risks associated with healing such as swelling, bleeding, possibility of additional surgeries, prolonged recovery, color and shape changes, infections, failure to meet the patient's goals and expectations, and added expense to the patient.

There may also be a longer recovery period due to the length of surgery and the anesthesia administered. Patients with significant skin laxity (like in a body lift procedure) will continue to have the same lax skin after surgery. The quality or elasticity of the skin will not change, and recurrence of skin looseness will occur at some time in the future, quicker for some than others. There are nerve endings that may become affected by healing scars from the surgery. While there may not be a major nerve injury, the small nerve endings may become too active during the healing period, producing a painful or oversensitive area due to their associations with the scar tissue. Often, massage and early non-surgical interventions can resolve this. It is important to discuss post-surgical pain with your surgeon.

Bleeding:

It is possible, though unusual, to experience a bleeding episode during or after the surgery. Should post-operative bleeding occur, it may require emergency treatment to drain accumulated blood, or you may require a blood transfusion, though such occurrences are rare. The collection of blood that can occur under your skin following surgery is referred to as a hematoma. Increased activity too soon after surgery can lead to increased chance of bleeding and additional surgeries. It is important to follow postoperative instructions and limit exercise and strenuous activity for the instructed time. Non-prescription "herbs" and dietary supplements can increase the risk of surgical bleeding. Hematoma can occur at any time, usually in the first three weeks following injury to the operative area. If blood transfusions are necessary to treat blood loss, there is the risk of blood-related infections such as hepatitis and HIV (AIDS). Your surgeon may provide medications after your surgery to prevent blood clots. Medications that are used to prevent blood clots in veins can produce bleeding and decreased blood platelets.

Infection:

Infection, although uncommon, can occur after surgery. Should an infection occur, additional treatment, including antibiotics, hospitalization, or even surgery, may be necessary. It is important to tell your surgeon of any other infections, such as a history of methicillin-resistant *Staphylococcus aureus* (MRSA) infections, an open wound, recent upper respiratory infection/pneumonia, ingrown toenail, insect bite, tooth abscess, or urinary tract infection. Infections in other parts of the body may lead to an infection in the operated area. Post-operative infections often result in more extensive scarring and predispose the patient to revision surgery.

Scarring:

All surgeries leave scars, some more visible than others. Although good wound healing after a surgical procedure is expected, abnormal scarring may occur within the skin and deeper tissues. Scars may be unattractive, and of a different color than the surrounding skin tone. Scar appearance may also vary within the same scar. Scars may be asymmetrical (appear different on the right and left side of the body). There is a possibility of visible marks in the skin from sutures. In some cases, scars may require surgical revision or treatment.

Firmness:

Excessive firmness can occur after surgery due to internal scarring. The occurrence of this is not predictable. Additional treatments including surgery may be necessary.

Patient Name:

Patient Initials: _____

Skin Sensitivity:

Itching, tenderness, or exaggerated responses to hot or cold temperatures may occur after surgery. This usually resolves during healing, but in rare situations, it may be chronic.

Major Wound Separation:

Wounds may separate after surgery. Should this occur, additional treatment, including surgery, may be necessary.

Sutures:

Most surgical techniques use deep sutures. You may notice these sutures after your surgery. Sutures may spontaneously poke through the skin, become visible, or produce irritation that requires suture removal.

Damage to Deeper Structures:

There is the potential for injury to deeper structures including nerves, blood vessels, muscles, and lungs (pneumothorax) during any surgical procedure. The potential for this to occur varies according to the type of procedure being performed. Injury to deeper structures may be temporary or permanent.

Fat Necrosis:

Fat tissue found deep in the skin might die. This may produce areas of firmness within the skin. Additional surgery to remove areas of fat necrosis may be necessary. There is a possibility of contour irregularities in the skin that may result from fat necrosis.

Surgical Anesthesia:

Both local and general anesthesia involve risks. There is a possibility of complications, injuries, and even death from all forms of surgical anesthesia or sedation.

Shock:

In rare circumstances, your surgical procedure can cause severe trauma, particularly when multiple or extensive procedures are performed. Although serious complications are infrequent, infections or excessive fluid loss can lead to severe illness and even death. If surgical shock occurs, hospitalization and additional treatment will be necessary.

Pain:

You will experience pain after your surgery. Pain of varying intensity and duration may occur and persist after surgery. If you are a chronic pain patient followed by a pain therapy practitioner, you may be asked to see this practitioner

preoperatively to assist you in the management of your pain disorder during the post-operative period. Chronic pain may occur infrequently from nerves becoming trapped in scar tissues or due to tissue stretching.

There are nerve endings that may be affected by healing scars from the surgery. While there may not be a major nerve injury, small nerve endings may become too active during the healing period, producing a painful or oversensitive area when they are involved with scar tissues. Often, massages and early non-surgical interventions can resolve this issue. It is important to discuss post-surgical pain with your surgeon.

Patient Name:

Patient Initials: _____

Cardiac and Pulmonary Complications:

Pulmonary complications may occur secondarily to blood clots (pulmonary emboli), fat deposits (fat emboli), pneumonia, or partial collapse of the lungs after general anesthesia. Pulmonary emboli can be life threatening or fatal in some circumstances. Inactivity and other conditions may increase the incidence of blood clots traveling to the lungs, causing a major blood clot that may result in death. It is important to discuss with your physician any past history of swelling in your legs or blood clots that may contribute to this condition. Cardiac complication is a common risk with any surgery and anesthesia, even in patients without symptoms. If you experience shortness of breath, chest

pains, or unusual heartbeats, seek medical attention immediately. Should any of these complications occur, you may require hospitalization and additional treatments.

Venous Thrombosis (Clot) and Sequelae:

Thrombosed veins, which resemble cords, occasionally develop in the area of the breast or around IV sites, and usually resolve without medical or surgical treatments. It is important to discuss with your surgeon any birth control pills you are taking. Certain high estrogen pills may increase your risk of thrombosed veins. Personal history of bleeding and clotting problems may also increase your risk of thrombosed veins.

Allergic Reactions:

In rare cases, local allergies to tape, suture material and glues, blood products, topical preparations, and injected agents have been reported. Serious systemic reactions including shock (anaphylaxis) may occur in response to drugs used during surgery and prescription medicines. Allergic reactions may require additional treatments. It is important to notify your physician of any previous allergic reactions.

Drug Reactions:

Unexpected drug allergies, lack of proper response to medication, or illness caused by prescribed drugs may occur. It is important for you to inform your physician of any problems and allergies you have had with any prescribed or over the counter medications, as well as medications you are currently taking on a regularly basis. Provide your surgeon with a list of medications and supplements you are currently taking.

Surgical Wetting Solutions:

There is a possibility that the large volumes of fluid containing dilute local anesthetic drugs and epinephrine injected into fat deposits during surgery may contribute to fluid overload or systemic reactions to these medications. Additional treatments including hospitalization may be necessary.

Persistent Swelling (Lymphedema):

Persistent swelling can occur following surgery.

Unsatisfactory Result:

Although good results are expected, there is no guarantee or warranty on the final results. The body is not symmetric, and almost everyone has some degree of unevenness, which may not be recognized in advance. One side of the face may be slightly larger, and one side of the face may be droopier. Similar possibilities exist for the breast and trunk areas. Many such issues cannot be fully corrected with surgery. The more realistic your expectations are, the better your results will appear to you. Some patients never achieve their desired goals or results, but at no fault of the surgeon or surgery. You may be disappointed with the results of surgery. Asymmetry, unanticipated shape and size, loss of function, wound disruption, poor healing, and loss of sensation may occur after surgery. Size may be incorrect. The location or appearance of surgical scar may be unsatisfactory. It may be necessary to perform additional surgeries to improve your results. Unsatisfactory results may NOT improve with each additional treatment.

Patient Name:

Patient Initials: _____

ADDITIONAL ADVISORIES

Medications and Herbal Dietary Supplements:

There are potential adverse reactions that occur as a result of taking over-the-counter, herbal, and/or prescription medications. Aspirin and medications that contain aspirin interfere with forming blood clots, and therefore may contribute to more bleeding issues. If you have a medical condition (such as heart arrhythmia, heart stent, blood vessels with blockages, or blood clots), and are taking anticoagulant medications such as Plavix®, Coumadin®, Xarelto®, Effient®, or Pradaxa® to thin your blood and prevent clotting, discuss management of these medications around the time of surgery with your plastic surgeon. Your plastic surgeon may sometimes coordinate a plan for these medications with the doctor that prescribed them for your medical condition. If you have been prescribed drugs for a medical condition, do not stop them without discussing it first with your plastic surgeon. Abruptly stopping these medications may result in heart attacks, strokes, or death. Be sure to check with your physician about any drug interactions that may exist with medications that you are already taking. If you have an adverse reaction, stop the drugs immediately and call your plastic surgeon for further instructions. If the reaction is severe, immediately go to the nearest emergency room.

When taking the prescribed pain medications after surgery, realize that they can affect your thought process and coordination. Do not drive, operate complex equipment, make any important decisions, or drink alcohol while taking these medications. Be sure to take your prescribed medication only as directed.

Sun Exposure - Direct or Tanning Salon:

The effects of the sun are damaging to the skin. Sun exposure to the treated areas may result in increased scarring, color changes, and poor healing. Patients who tan, either outdoors or in a salon, should inform their surgeon and either delay treatment, or avoid tanning until the surgeon says it is safe to resume. The damaging effect of sun exposure occurs even with the use of sun block or clothing coverage.

Travel Plans:

Any surgery holds the risk of complications that may delay healing and your return to normal life. Please let the surgeon know of any travel plans, important commitments that were already scheduled or planned, or time demands that are important to you, so that surgery can occur at appropriate times. There are no guarantees that you will be able to resume all activities in the desired timeframe. Allow at least 10-14 days prior to travel via airplane. Medications may be required should you have a long flight/trip to prevent DVT/PE in the immediate post-operative period.

Long-term Results:

Subsequent alterations in the appearance of your body may occur as the result of aging, sun exposure, weight loss, weight gain, pregnancy, menopause or other circumstances not related to your surgery.

Body Piercing:

Individuals who currently wear body-piercing jewelry in the surgical region are advised that an infection could develop from this activity. Body-piercing jewelry should be removed prior to your surgical procedure.

Nails:

To determine your vitals status during surgery, your anesthesia provider may require access to your fingernails for monitoring. Make sure to have at least two fingernails free of nail polish or acrylic nails on the date of your surgery.

Jewelry:

Jewelry should not be brought with you at the time of your surgical procedure. Items, such as earrings, wedding rings, and necklaces should be removed and placed in a safe place.

Patient Name:

Patient Initials: _____

Future Pregnancy and Breastfeeding:

This surgery is not known to interfere with pregnancy. If you are planning a pregnancy, your breast skin may stretch and offset the results of surgery. You may have more difficulty breast-feeding after this operation.

Female Patient Information:

It is important to inform your plastic surgeon if you use birth control pills or estrogen replacement, or if you suspect you may be pregnant. Many medications including antibiotics may neutralize the preventive effect of birth control pills, allowing for conception and pregnancy.

Intimate Relations After Surgery:

Recovery from surgery involves coagulation of blood vessels, and increased activity of any kind may open these vessels, leading to bleeding, or hematoma. Activity that increases your pulse or heart rate may cause additional bruising, swelling, and the need for return to surgery to control bleeding. It is wise to refrain from intimate physical activities until your physician states it is safe.

Mental Health Disorders and Elective Surgery:

It is important that all patients seeking to undergo elective surgery have realistic expectations that focus on improvements rather than perfection. Complications or less-than-satisfactory results are sometimes unavoidable, may require additional surgeries, and are often stressful. Please openly discuss with your surgeon, prior to surgery, any past history of significant emotional depression or mental health disorders. Although many individuals may benefit psychologically from the results of elective surgery, effects on mental health cannot be accurately predicted.

ADDITIONAL SURGERY NECESSARY (Re-Operations)

There are many variable conditions that may influence the long-term result of the surgery. It is unknown how your tissue may respond or how wound healing will occur after surgery. Secondary surgeries may be necessary to perform additional tightening or repositioning of body structures. Should complications occur, additional surgery or other treatments may be necessary. Even though risks and complications occur infrequently, the risks cited above are associated with this surgery. Other complications and risks can occur but are less common. The practice of medicine and surgery is not an exact science. Although good results are expected, there is no guarantee or warranty on the expected results. In some situations, it may not be possible to achieve optimal results with a single surgical procedure. You and your surgeon will discuss the options available should additional surgeries be advised. There may be additional costs and expenses for such additional procedures, including surgical fees, facility and anesthesia fees, and pathology and lab testing.

PATIENT COMPLIANCE

Follow all physician's instructions carefully; this is essential for the success of your surgical outcome. It is important that the surgical incisions are not subjected to excessive force, swelling, abrasion, or motion during the time of healing. Personal and vocational activities need to be restricted. Protective dressings and drains should not be removed unless instructed by your plastic surgeon. Successful post-operative function depends on both surgery and subsequent care. Physical activities that increase your pulse or heart rate may cause bruising, swelling, fluid accumulation, and the need for return to surgery. It is important that you participate in follow-up care and return for aftercare to promote your recovery after surgery.

Patient Name:

Patient Initials: _____

ATTESTATIONS

Smoking, Second-Hand Smoke Exposure, Nicotine Products (Patch, Gum, Nasal Spray):

Patients who are currently smoking or use tobacco or nicotine products (patch, gum, or nasal spray) are at greater risk for significant surgical complications of skin loss, delayed healing, and additional scarring. Individuals exposed to second-hand smoke are also at potential risk for similar complications attributable to nicotine exposure. Additionally, smoking may have a significant negative effect on anesthesia and recovery from anesthesia, and can lead to coughing and possibly increased bleeding. Individuals who are not exposed to tobacco smoke or nicotine-containing products have a significantly lower risk of these types of complications. Please indicate your current status regarding these items below:

___ I am a non-smoker and do not use nicotine products. I understand the potential risk of second-hand smoke exposure causing surgical complications.

___ I am a smoker or use tobacco/nicotine products. I understand the risk of surgical complications due to smoking or use of nicotine products.

___ I have smoked and stopped approximately _____ ago. I understand I may still have the effects and therefore risks from smoking in my system, if not enough time has lapsed.

___ I have been advised to stop smoking immediately and have been informed of the risks, benefits, expectations, and alternatives to my surgery if I continue smoking.

It is important to refrain from smoking at least 6 weeks before surgery and until your physician states it is safe to return, if desired. I acknowledge that I will inform my physician if I continue to smoke within this timeframe, and understand that for my safety, the surgery, if possible, may be delayed.

Smoking may have such a negative effect on your surgery that a urine or blood test just before surgery may be done just before surgery to determine the presence of nicotine. If positive, your surgery may be cancelled. Your surgery, the scheduling fee, and other prepaid amounts may be forfeited. Honestly disclose smoking status to your surgeon.

PROCEED TO NEXT PAGE

Patient Name:

Patient Initials: _____

Sleep Apnea/CPAP:

Individuals who have breathing disorders such as “obstructive sleep apnea,” and who may rely upon CPAP devices (continuous positive airway pressure) or utilize nighttime oxygen are advised that they are at a substantive risk for respiratory arrest and death when they take narcotic pain medications following surgery. This is an important consideration when evaluating the safety of surgical procedures in terms of very serious complications, including death, that relate to pre-existing medical conditions. Surgery may be considered only with post-surgery monitoring is conducted in a hospital setting in order to reduce the risk of potential respiratory complications, and to safely manage pain following surgery.

Please consider the following symptoms of sleep apnea:

- I am frequently tired upon waking, and throughout the day
- I have trouble staying asleep at night
- I have been told that I snore or stop breathing during sleep
- I wake up throughout the night or constantly turn from side to side
- I have been told that my legs or arms jerk while I’m sleeping
- I make abrupt snorting noises during sleep
- I feel tired or fall asleep during the day

It is important for you to inform and discuss any of the above symptoms that you have experienced with your surgeon.

DVT/PE Risks and Advisory:

There is a risk of blood clots, DVT, and PE with every surgical procedure. It varies with the risk factors below. The higher the risk factors, the greater the risk, and the more involved you must be in both understanding these risks and, when permitted by your physician, walking and moving your legs. There may also be leg stockings, squeezing active leg devices, and possibly medicines to help lower your risk.

There are many conditions that may increase or affect risks of clotting. Inform your doctor about any past or present history of any of the following:

- Past History of Blood Clots
- Family History of Blood Clots
- Birth Control Pills
- Hormone Stimulating Drugs
- Swollen Legs
- History of Cancer
- Large Dose of Vitamins
- Varicose Veins
- Past Illnesses of the Heart, Liver, Lung, or Gastrointestinal Tract.
- History of Multiple Spontaneous Abortions or Miscarriages
- I understand the risks relating to DVT/PE, and how important it is to comply with therapy as discussed with my surgeon. The methods of preventative therapy include:
 - Early ambulation when allows
 - Compression devices (SCD/ICD)
 - Anticoagulation protocols when Allowed

For high risk patients, the risks of VTE are still high even with appropriate chemoprophylaxis. If your surgery is elective and you are a high risk patient, it is best to consider with not proceeding with such elective surgery.

Patient Name:

Patient Initials: _____

CONSENT for SURGERY/PROCEDURE or TREATMENT

1. I hereby authorize Dr. Michael Salzhauer and assistants who may be selected to perform **Rhinoplasty Surgery**.

I have received the following information sheet: **Rhinoplasty Surgery**.

2. I recognize that during the course of the operation and medical treatment or anesthesia, unforeseen conditions may necessitate different procedures than those outlined above. I therefore authorize the above physician and assistants or designees to perform such other procedures, which are deemed necessary and desirable, based on his or her professional judgment. The authority granted under this paragraph shall include all conditions that require treatments and those not known to my physician at the time the procedure has begun.

3. I consent to the administration of such anesthetics as considered necessary or advisable. I understand that all forms of anesthesia involve risks and the possibility of complications, injury, and sometimes death.

4. I understand what my surgeon can and cannot do, and understand that there are no warranties or guarantees, implied or specific about my outcome. I have had the opportunity to explain my goals, and understand which desired outcomes are realistic and which are not. All of my questions have been answered, and I understand the inherent (specific) risks to the procedures I seek, as well as those additional risks and complications, benefits, and alternatives. Understanding all of this, I elect to proceed.

5. I consent to be photographed or televised before, during, and after the operation(s) or procedure(s) to be performed, including appropriate portions of my body, for medical, scientific, or educational purposes, provided my identity is not revealed by the pictures.

6. For purposes of advancing medical education, I consent to the admittance of observers to the operating room.

7. I consent to the disposal of any tissue, medical devices, or body parts that may be removed.

8. I am aware that there are potential significant risks to my health with the utilization of blood products, and I consent to their utilization should they be deemed necessary by my surgeon and/or his/her appointees.

9. I authorize the release of my Social Security number to appropriate agencies for legal reporting and medical-device registration, if applicable.

10. I understand that the surgeons' fees are separate from the anesthesia and hospital charges, and the fees are agreeable to me. If a secondary procedure is necessary, further expenditure will be required.

11. I realize that not having the operation is an option. I opt out of having this procedure ____.

12. IT HAS BEEN EXPLAINED TO ME IN A WAY THAT I UNDERSTAND:

- a. THE ABOVE TREATMENT OR PROCEDURE TO BE UNDERTAKEN
- b. THERE MAY BE ALTERNATIVE PROCEDURES OR METHODS OF TREATMENT
- c. THERE ARE RISKS TO THE PROCEDURE OR TREATMENT PROPOSED

I CONSENT TO THE TREATMENT OR PROCEDURE AND THE ABOVE LISTED ITEMS (1-12). I AM SATISFIED WITH THE EXPLANATION.

Patient:

(or Person Authorized to Sign for Patient)

Date:

Witness Signature:

Date:

Physician Signature:

Date: